arXiv:1407.8496v2 [math.CO] 14 Apr 2015

On association schemes with at most two nonlinear irreducible characters

Javad Bagherian
Department of Mathematics, University of Isfahan,
Isfahan 81746-73441, Iran.
bagherian@sci.ui.ac.ir

April 15, 2015

Abstract

An irreducible character χ of an association scheme (X, S) is called nonlinear if the multiplicity of χ is greater than 1. The main result of this paper gives a characterization of commutative association schemes with at most two nonlinear irreducible characters. This yields a characterization of finite groups with at most two nonlinear irreducible characters. A class of noncommutative association schemes with at most two nonlinear irreducible character is also given.

Key words : association scheme, nonlinear character, group-like scheme.
AMS Classification: 05E30.

1 Introduction

In the character theory of association schemes, the character values of irreducible characters give many useful information about association schemes. In particular, the multiplicities of irreducible characters paly an important role for the determining the structure of association schemes. An interesting problem about the multiplicities of irreducible characters is what can be said about (X, S) when the number of nonlinear irreducible characters is known. An irreducible character χ of an association scheme (X, S) is called nonlinear if $m_\chi > 1$. In this paper we characterize the structure of commutative association schemes with at most two nonlinear irreducible characters. This yields a characterization of finite groups with at most two nonlinear irreducible characters. Moreover, we give a characterization of noncommutative association schemes with one nonlinear irreducible characters. Finally, a class of noncommutative association schemes with two nonlinear irreducible characters is given.

2 Preliminaries

Let us first state some necessary definitions and notation. For details, we refer the reader to [11, 12] for the background of association schemes. Throughout this paper, $\mathbb{C}$ denotes the complex numbers.

2.1 Association schemes

Definition 2.1. *Let X be a finite set and $S = \{s_0, s_1, \ldots, s_n\}$ be a partition of $X \times X$. Then (X, S) is called an association scheme with n classes if the following properties hold:*

(i) $s_0 = \{(x, x) | x \in X\}$.

(ii) *For every $s \in S$, s^* is also in S, where $s^* := \{(x, y) | (y, x) \in s\}$.*

(iii) *For every $g, h, k \in S$, there exists a nonnegative integer λ_{ghk} such that for every $(x, y) \in k$, there exist exactly λ_{ghk} elements $z \in X$ with $(x, z) \in g$ and $(z, y) \in h$.*

The diagonal relation s_0 will be denoted by 1. For each $s \in S$, we call $n_s = \lambda_{ss^*1}$ the *valency* of s. For any nonempty subset H of S, put $n_H = \sum_{h \in H} n_h$. We call n_S the *order* of (X, S). An association scheme (X, S) is called *commutative* if for all $g, h, k \in S$, $\lambda_{ghk} = \lambda_{hgk}$. Let H and K be nonempty subsets of S. We define HK to be the set of all elements $t \in S$ such that there exist element $h \in H$ and $k \in K$ with $\lambda_{hkt} \neq 0$. The set HK is called the *complex product* of H and K. If one of factors in a complex product consists of a single element s, then one usually writes s for $\{s\}$. A nonempty subset H of S is called a *closed subset* if $HH \subseteq H$. A closed subset H of S is called *strongly normal*, denoted by $H \lhd^\sharp S$, if $sHs^* = H$ for any $s \in S$. We put $\mathrm{O}^\vartheta(S) = \bigcap_{H \lhd^\sharp S} H$ and call it the *thin residue* of H. For a closed subset H of S, the number n_S/n_H is called the *index* of H in S and is denoted by $|S : H|$.

Let (X, S) be a scheme and H be a closed subset of S. For every $x \in X$ put $xH = \bigcup_{h \in H} xh$, where $xh = \{y \in X | (x, y) \in h\}$. A *subscheme* $(X, S)_{xH}$ is a scheme with the points xH and the set of relations $\{s_{xH} | s \in H\}$, where $s_{xH} = s \bigcap (xH \times xH)$.

Moreover, if we put $X/\!\!/H = \{xH | x \in X\}$ and $S/\!\!/H = \{s^H | s \in S\}$ where $s^H = \{(xH, yH) | y \in xHsH\}$, then $(X/\!\!/H, S/\!\!/H)$ is a scheme, called the *quotient scheme* of (X, S) over H. Note that a closed subset H is strongly normal if and only if the quotient scheme $(X/\!\!/H, S/\!\!/H)$ is a group with respect to the relational product if and only if $ss^* \subseteq H$, for every $s \in S$ (see [11, Theorem 2.2.3]). In particular, since $S/\!\!/\mathrm{O}^\vartheta(S)$ is a finite group, we can consider the derived subgroup $(S/\!\!/\mathrm{O}^\vartheta(S))'$ of $S/\!\!/\mathrm{O}^\vartheta(S)$. Suppose S' be the inverse image of $(S/\!\!/\mathrm{O}^\vartheta(S))'$. Then quotient scheme $(X/\!\!/S', S/\!\!/S')$ is a abelian group with respect to the relational product.

2.2 Characters of association schemes

Let (X, S) be an association scheme. For every $s \in S$, let σ_s be the adjacency matrix of s. For convenience, σ_1 is denoted by 1. It is known that $\mathbb{C}S = \bigoplus_{s\in S} \mathbb{C}\sigma_s$, the *adjacency algebra* of (X, S), is a semisimple algebra (see [11, Theorem 4.1.3]). The set of irreducible characters of S is denoted by $\mathrm{Irr}(S)$. One can see that $1_S \in \mathrm{Hom}_{\mathbb{C}}(\mathbb{C}\mathrm{S}, \mathbb{C})$ such that $1_S(\sigma_s) = n_s$ is an irreducible character of $\mathbb{C}S$, called the *principal* character. In [5], Hanaki has shown that the irreducible characters of $S/\!\!/\mathrm{O}^{\vartheta}(S)$ can be considered as irreducible characters of S.

Let Γ_S be a representation of $\mathbb{C}S$ which sends σ_s to itself for every $s \in S$. Let γ_S be the character afforded by Γ_S. Then one can see that $\gamma_S(1) = |X|$ and $\gamma_S(\sigma_s) = 0$ for every $1 \neq s \in S$. Consider the following irreducible decomposition of γ_S,

$$\gamma_S = \sum_{\chi\in\mathrm{Irr}(S)} m_\chi \chi.$$

We call m_χ the *multiplicity* of χ. One can see that $m_{1_S} = 1$ and $|X| = \sum_{\chi\in\mathrm{Irr}(S)} m_\chi \chi(1)$ (see [12, section 4]).

Example 2.2. *Let G be a finite group and $C_0 = \{1\}, C_1, \ldots, C_h$ be the conjugacy classes of G. Define R_i by $(x, y) \in R_i$ if and only if $xy^{-1} \in C_i$, where $x, y \in G$. Put $S = \{R_i\}_{0\leq i\leq h}$. Then (G, S) is an association scheme, which is called the group association scheme of G. For every relation R_i of S, $n_{R_i} = |C_i|$ and one can see that $R_i \in \mathrm{O}^{\vartheta}(S)$ if and only if $C_i \subseteq G'$, where G' is the derived subgroup of G.*

The adjacency algebra of (G, S) is isomorphic to the algebra $\mathrm{Z}(\mathbb{C}G)$ with the basis $\mathrm{Cla}(G)$, where $\mathrm{Z}(\mathbb{C}G)$ is the center of group algebra $\mathbb{C}G$ and $\mathrm{Cla}(G) = \{K_0, \cdots, K_h\}$, with $K_i = \sum_{g\in C_i} g$. Furthermore, $\{\omega_\chi |\ \chi \in \mathrm{Irr}(G)\}$ is the set of irreducible characters of $Z(\mathbb{C}G)$, where

$$\omega_\chi(K_i) = \frac{\chi(g)|C_i|}{\chi(1)},$$

for some $g \in C_i$. Since $|G| = \sum_{\chi\in\mathrm{Irr}(G)} \chi(1)^2$, it follows that $m_{\omega_\chi} = \chi(1)^2$, for every $\chi \in \mathrm{Irr}(G)$.

Let (X, S) be an association scheme and T be a closed subset of S. Suppose that L is a $\mathbb{C}T$-module which affords the character φ, and V is a $\mathbb{C}S$-module which affords the character χ. Then V is a $\mathbb{C}T$-module which affords the restriction χ_T of χ to $\mathbb{C}T$, and $L^S = L \otimes_{\mathbb{C}T} \mathbb{C}S$ is a $\mathbb{C}S$-module which affords the induction φ^S of φ. Suppose that T is strongly normal. Put $G = S/\!\!/T$. Let φ be an irreducible character of T and L be an irreducible $\mathbb{C}T$ module affording φ. Consider the induction of L to S. Then

$$L^S = L \otimes_{\mathbb{C}T} \mathbb{C}S = \bigoplus_{s^T\in S/\!\!/T} L \otimes \mathbb{C}(TsT).$$

The *stabilizer* $G\{L\}$ of L in G is defined by

$$G\{L\} = \{s^T \in S/\!\!/T | L \otimes \mathbb{C}(TsT) \cong L\}.$$

One can see that $G\{L\}$ is a subgroup of G. The set of $S/\!\!/T$-conjugates of L is $\{L \otimes \mathbb{C}(TsT) | s \in S,\ L \otimes \mathbb{C}(TsT) \neq 0\}$. From [1] it follows that if L and L' are $S/\!\!/T$-conjugates, then $L^S \cong L'^S$.

Theorem 2.3. *(See [1].) Let (X, S) be a scheme and T be a strongly closed subset of S. Put $G = S/\!\!/T$. Then for every $\chi \in \mathrm{Irr}(S)$, there exists a positive integer e such that*

$$\chi_T = e \sum_{i=1}^{n} \varphi_i,$$

where $\varphi_i, 1 \leq i \leq n$, are $S/\!\!/T$-conjugate irreducible characters of T.

Theorem 2.4. *(See [3].) Let (X, S) be an association scheme and T be a strongly normal closed subset of S. Suppose that $G = S/\!\!/T$ is the cyclic group of prime order p. Suppose that $\mathrm{Irr}(G) = \{\zeta_i | 1 \leq i \leq p\}$. Then for $\chi \in \mathrm{Irr}(S)$, one of the following statements holds:*

(1) *$\chi_T \in \mathrm{Irr}(T)$ and $(\chi_T)^S = \sum_{i=1}^{p} \chi\zeta_i$,*

(2) *$\chi(\sigma_s) = 0$, for any $s \in S \setminus T$ and χ_T is a sum of at most p distinct irreducible characters. If ψ is an irreducible constituent of χ_T, then $\psi^S = \chi$.*

The product of characters in schemes has been given in [4]. If (X, S) is a scheme and T is a strongly normal closed subset of S, then it follows from [4, Theorem 3.3] that for every $\chi \in \mathrm{Irr}(S)$ and $\zeta \in \mathrm{Irr}(S/\!\!/T)$, the character product $\chi\zeta$ defined by

$$\chi\zeta(\sigma_s) = \chi(\sigma_s)\zeta(\sigma_{s^T})$$

is an irreducible character of S. Then $\mathrm{Irr}(S/\!\!/T)$ acts on $\mathrm{Irr}(S)$ by the above multiplication.

2.3 Wreath product of association schemes

Let (X, S) and (Y, F) be association schemes. For every $s \in S$ we define

$$\overline{s} = \{((x_1, y), (x_2, y)) | y \in Y, (x_1, x_2) \in s\}.$$

For every $f \in F \setminus \{1_Y\}$ we define

$$\overline{f} = \{((x_1, y_1), (x_2, y_2)) | x_1, x_2 \in X, (y_1, y_2) \in h\}.$$

Then $(X \times Y, S \wr F)$ is an association scheme called the *wreath product* of (X, S) and (Y, F), where

$$S \wr F = \{\overline{s} | s \in S\} \cup \{\overline{f} | f \in F \setminus \{1_Y\}\}.$$

Lemma 2.5. *(See [10, Lemma 3.1].) Let (X, S) be a commutative association scheme and H be a closed subset of S. Then the following are equivalent:*

(1) *S is the wrath product of association schemes $(X, S)_{xH}$ and $(X/\!\!/H, S/\!\!/H)$,*

(2) *$|S| = |H| + |S/\!\!/H| - 1$,*

(3) *For every $h \in H$ and $s \in S \setminus H$, $\sigma_s\sigma_h = \sigma_h\sigma_s = n_h\sigma_s$.*

The following easy lemma is useful.

Lemma 2.6. *Let (X, S) be a commutative association scheme and H be a strongly normal closed subset of S. If S is the wrath product of association schemes $(X, S)_{xH}$ and $(X/\!\!/H, S/\!\!/H)$, then $n_s = n_H$ for every $s \in S \setminus H$.*

Proof. For every $s \in S \setminus H$ we have

$$n_{s^H} n_H = n_{Hs},$$

(see [11, Theorem 1.5.4(v)]). Since $n_{s^H} = 1$ so $n_H = n_{Hs}$. On the other hand, it follows from Lemma 2.5, that $Hs = \{s\}$. Hence $n_s = n_H$.

■

Let $\chi_1 = 1_S, \ldots, \chi_r$ be all irreducible characters of (X, S) and let $\psi_1 = 1_F, \ldots, \psi_t$ be all irreducible characters of (Y, F). We define

$$\begin{cases} \widetilde{\chi}_i(\sigma_{\overline{s}}) = \chi_i(\sigma_s), \\ \widetilde{\chi}_i(\sigma_{\overline{f}}) = 0, \end{cases}$$

for $i \neq 1$ and

$$\begin{cases} \widetilde{\psi}_j(\sigma_{\overline{s}}) = n_s\psi_j(1_Y), \\ \widetilde{\psi}_j(\sigma_{\overline{f}}) = n_S\psi(\sigma_f). \end{cases}$$

Then we have the following result.

Theorem 2.7. *(see [6]) In the above notation, $\widetilde{\chi_2}, \ldots, \widetilde{\chi}_r, \widetilde{\psi_1}, \ldots, \widetilde{\psi}_t$ are all irreducible characters of $(X \times Y, S \wr F)$.*

2.4 Group-like schemes

Let (X, S) be an association scheme. We define a binary relation $\sim$ on S as follows. For $s, t \in S$, we write $s \sim t$ if

$$\chi(\sigma_s)/n_s = \chi(\sigma_t)/n_t, \tag{1}$$

for every $\chi \in \mathrm{Irr}(S)$. Then $\sim$ is an equivalence relation. For $s \in S$, put $\widetilde{s} = \bigcup_{t \sim s} s$ and $\widetilde{S} = \{\widetilde{s} | s \in S\}$. If $Z(\mathbb{C}S) = \bigoplus_{\widetilde{s} \in \widetilde{S}} \mathbb{C}\sigma_{\widetilde{s}}$, then (X, S) is called a *group-like scheme*.

Theorem 2.8. *(See [2, Theorem 4.1].) For an association scheme (X, S), the following statements are equivalent:*

(1) *(X, S) is a group-like scheme,*

(2) $\dim(\mathrm{Z}_{\mathbb{C}}(\mathbb{C}\mathrm{S}) = |\widetilde{\mathrm{S}}|$,

(3) *for every $\chi, \psi \in \mathrm{Irr}(S)$, $\chi\psi$ is a linear combination of* $\mathrm{Irr}(S)$.

The following easy lemma is useful.

Lemma 2.9. *An association scheme (X, S) is group-like scheme if and only if for every $\chi, \psi \in \mathrm{Irr}(S)$ and every $s, h \in S$ we have $\chi\psi(\sigma_s\sigma_h) = \chi\psi(\sigma_h\sigma_s)$.*

Proof. For every $\chi, \psi \in \mathrm{Irr}(S)$, consider the linear function $\chi\psi : \mathbb{C}S \to \mathbb{C}$ by

$$\chi\psi(\sum_{s\in S} \lambda_s\sigma_s) = \sum_{s\in S} \lambda_s\chi\psi(\sigma_s).$$

It follows from [4, Theorem 4.2] that $\chi\psi$ is a linear combination of irreducible characters if and only if $\chi\psi(\sigma_s\sigma_h) = \chi\psi(\sigma_h\sigma_s)$ for every $s, h \in S$. The result now follows from Theorem 2.8.

■

3 Main Results

Let G be a finite group. An irreducible character χ of G is called nonlinear if $\chi(1) > 1$. In this section we first define the concept of the nonlinear irreducible character for association schemes and then give a characterization of association schemes with at most two nonlinear irreducible character.

Let (X, S) be an association schemes. We say that an irreducible character χ of S is *linear* if $m_\chi = 1$; otherwise χ is called *nonlinear*. It follows from [8, Lemma 2.4(v)] that $\chi \in \mathrm{Irr}(S/\!\!/S')$ if and only if $m_\chi = 1$. So $|S : S'|$ is the number of linear characters of S and $\mathrm{Irr}(S) - \mathrm{Irr}(S/\!\!/S')$ is the set of nonlinear characters of S. In particular, if (X, S) is commutative, then an irreducible character $\chi \in \mathrm{Irr}(S)$ is linear if and only if $\chi \in \mathrm{Irr}(S/\!\!/\mathrm{O}^\vartheta(S))$ and so $\mathrm{Irr}(S) - \mathrm{Irr}(S/\!\!/\mathrm{O}^\vartheta(S))$ is the set of nonlinear characters of S.

If (G, S) is the group association scheme of G, then an irreducible character χ of G is nonlinear if and only if the irreducible character ω_χ of S is nonlinear; see Example 2.2.

3.1 Association schemes with one nonlinear irreducible character

In this section we give a characterization of association schemes with exactly one nonlinear irreducible character.

Lemma 3.1. *A commutative association scheme (X, S) has exactly one nonlinear irreducible character if and only if $|\mathrm{O}^{\vartheta}(S)| = 2$ and S is the wreath product of association schemes $(X, S)_{x\mathrm{O}^{\vartheta}(S)}$ and $(X/\!\!/\mathrm{O}^{\vartheta}(S), S/\!\!/\mathrm{O}^{\vartheta}(S))$ for $x \in X$.*

Proof. First we assume that S contains exactly one nonlinear irreducible character. Then $|S| = |S/\!\!/\mathrm{O}^{\vartheta}(S)| + 1$. Put $T = \mathrm{O}^{\vartheta}(S)$. Suppose that $S/\!\!/T = \{1^T, s_1^T, \ldots, s_n^T\}$, for some relations $s_i \in S$. Since $|S| = |S/\!\!/T| + 1$, it follows that $S = \{1, t, s_1, \ldots, s_n\}$ and $T = \{1, t\}$, for some relation t. Now since

$$|S| = |S/\!\!/T| + 1 = |T| + |S/\!\!/T| - 1,$$

it follows from Lemma 2.5 that S is the wreath product of association schemes $(X, S)_{xT}$ and $(X/\!\!/T, S/\!\!/T)$ for some $x \in X$.

Conversely, it follows from Theorem 2.7 that S contains exactly one nonlinear irreducible character.

■

From above lemma we can give the following characterization of finite groups with exactly one nonlinear irreducible character.

Corollary 3.2. *A finite group G has exactly one nonlinear irreducible character if and only if the derived subgroup G' of G contains two conjugacy classes and for every $g \in G \setminus G'$, coset gG' is the conjugacy class of G containing g.*

Proof. Let (G, S) be the group association scheme of G. Suppose $\{C_0 = \{1\}, \ldots, C_d\}$ is the conjugacy classes of G. Then we can consider $\{K_0, \ldots, K_d\}$ as a basis for the complex adjacency algebra of (G, S), where $K_i = \sum_{g \in C} g$; see Example 2.2. First assume that G has exactly one nonlinear irreducible character. Then (G, S) also has one nonlinear irreducible character and it follows from Lemma 3.1 that $|\mathrm{O}^{\vartheta}(S)| = 2$ and S is the wreath product of association schemes $(G, S)_{g\mathrm{O}^{\vartheta}(S)}$ and $(G/\!\!/\mathrm{O}^{\vartheta}(S), S/\!\!/\mathrm{O}^{\vartheta}(S))$ for $g \in G$. This implies that G' contains two conjugacy classes C_0 and C_i for some $1 \leq i \leq d$ and Lemma 2.5 shows that $K_iK_j = K_jK_i = |C_i|K_j$ for every $j \neq 0, i$. So we conclude that $C_iC_j = C_j$ for every $j \neq 0, i$. Hence for every $g \in C_j$, $gC_i \subseteq C_j$. Now from Lemma 2.6 we have $|C_j| = |G'| = |C_i| + 1$ and so $gG' = C_j$.

Conversely, since G' contains two conjugacy classes C_0 and C, we have $|\mathrm{O}^{\vartheta}(S)| = 2$ and moreover, since for every conjugacy class $C' \notin \{C_0, C\}$ and $g \in C'$, $C' = gG'$ we get $C'C = C'$, and so $K'K = KK' = |C|K'$. Then it follows from Lemma 2.5 that S is the wreath product of association schemes $(G, S)_{g\mathrm{O}^{\vartheta}(S)}$ and $(X/\!\!/\mathrm{O}^{\vartheta}(S), S/\!\!/\mathrm{O}^{\vartheta}(S))$ for $g \in G$. So Lemma 3.1 shows that (G, S) contains exactly one nonlinear irreducible character and hance G has exactly one nonlinear irreducible character.

■

Definition 3.3. *(See [9].) A finite group G is called Con-Cos group if G' contains two conjugacy classes and for every $g \in G \setminus G'$, coset gG' is the conjugate class containing g.*

Corollary 3.4. *A finite group G has one nonlinear irreducible character if and only if G is a Con-Cos group.*

Proof. The result follows immediately from Corollary 3.2.

■

Theorem 3.5. *An association scheme (X, S) has exactly one nonlinear irreducible character if and only if it is a group-like scheme, $|\mathrm{O}^\vartheta(\widetilde{S})| = 2$ and $\widetilde{S}$ is the wreath product of association schemes $(X, \widetilde{S})_{x\mathrm{O}^\vartheta(\widetilde{S})}$ and $(X/\!\!/\mathrm{O}^\vartheta(\widetilde{S}), S/\!\!/\mathrm{O}^\vartheta(\widetilde{S}))$ for $x \in X$.*

Proof. Suppose that (X, S) has exactly one nonlinear irreducible character. We first note that for every $s, t \in S \setminus S'$, we have $s^{S'} = t^{S'}$ if and only if for every $\psi \in \mathrm{Irr}(S/\!\!/S')$,

$$\psi(\sigma_s)/n_s = \psi(\sigma_{s^{S'}}) = \psi(\sigma_{t^{S'}}) = \psi(\sigma_t)/n_t$$

(see [5, Theorem 3.5]). On the other hand, since $S/\!\!/S'$ is an abelian group, $\mathrm{Irr}(S/\!\!/S')$ acts on $\mathrm{Irr}(S)$ by

$$\psi\zeta(\sigma_s) = \psi(\sigma_s)\zeta(\sigma_{s^{S'}})$$

for every $\psi \in \mathrm{Irr}(S)$ and $\zeta \in \mathrm{Irr}(S/\!\!/S')$; (see [4, Theorem 3.4]). Since χ is the only irreducible character with $m_\chi > 1$ we conclude that the orbit of χ has length 1. This implies that for every $\zeta \in \mathrm{Irr}(S/\!\!/S')$, $\chi\zeta = \chi$. Since for every $s \in S \setminus S'$, there exists $\zeta \in \mathrm{Irr}(S/\!\!/S')$ such that $\zeta(\sigma_{s^{S'}}) \neq 1$, from equality $\chi(\sigma_s) = \chi(\sigma_s)\zeta(\sigma_{s^{S'}})$ we have $\chi(\sigma_s) = 0$. Then for every $s, t \in S \setminus S'$, we have $s^{S'} = t^{S'}$ if and only if for every $\psi \in \mathrm{Irr}(S)$, $\psi(\sigma_s)/n_s = \psi(\sigma_t)/n_t$.

Now let $1 \neq s \in S'$. Since $\sum_{\psi \in \mathrm{Irr}(S)} m_\psi \psi(\sigma_s) = 0$, it follows that

$$m_\chi \chi(\sigma_s) + \sum_{\chi \neq \psi \in \mathrm{Irr}(S)} m_\psi \psi(\sigma_s) = 0.$$

Then we have

$$\chi(\sigma_s) = \frac{-|S/\!\!/S'|n_s}{m_\chi}$$

and hence

$$\chi(\sigma_s)/n_s = \frac{-|S/\!\!/S'|}{m_\chi}. \tag{2}$$

So for every $t \in S'$, (2) shows that

$$\frac{\chi(\sigma_t)}{n_t} = \frac{-|S/\!\!/S'|}{m_\chi} = \frac{\chi(\sigma_s)}{n_s}.$$

Since for every $\psi \in \mathrm{Irr}(S/\!\!/S')$ we have

$$\frac{\psi(\sigma_s)}{n_s} = \frac{\psi(\sigma_t)}{n_t} = 1,$$

it follows that

$$\frac{\psi(\sigma_s)}{n_s} = \frac{\psi(\sigma_t)}{n_t},$$

for every $\psi \in \mathrm{Irr}(S)$.

Now let $\sim$ be the equivalence relation which is defined in (1). From above we conclude that

$$\widetilde{S} = 1 \cup \widetilde{t} \cup \{s^{S'} | s \in S \setminus S'\},$$

for some $t \in S'$. Then $|\widetilde{S}| = |S/\!\!/S'| + 1$. Since $\dim_{\mathbb{C}}\mathrm{Z}(\mathbb{C}\mathrm{S}) = |\mathrm{Irr(S)}| = |\mathrm{S}/\!\!/\mathrm{S}'| + 1$ we have $\mathbb{C}\widetilde{S} = Z(\mathbb{C}S)$ and it follows from Theorem 2.8 that (X, S) is a group-like scheme. Since $(X, \widetilde{S})$ has exactly one nonlinear irreducible character, Lemma 3.1 shows that $|\mathrm{O}^{\vartheta}(\widetilde{S})| = 2$ and $\widetilde{S}$ is the wreath product of association schemes $(X, \widetilde{S})_{x\mathrm{O}^{\vartheta}(\widetilde{S})}$ and $(X/\!\!/\mathrm{O}^{\vartheta}(\widetilde{S}), S/\!\!/\mathrm{O}^{\vartheta}(\widetilde{S}))$ for $x \in X$.

Conversely, it follows from Lemma 3.1 that $(X, \widetilde{S})$ has exactly one nonlinear irreducible character. Since $m_{\widetilde{\chi}} = \chi(1)m_{\chi}$, for every $\chi \in \mathrm{Irr}(S)$ we conclude that (X, S) has exactly one nonlinear irreducible character. The proof is now complete. ∎

3.2 Association schemes with two nonlinear irreducible characters

In this section we first give a characterization of commutative association schemes with exactly two nonlinear irreducible characters. Then we give a class of association schemes with two nonlinear irreducible characters.

Theorem 3.6. *Let (X, S) be a commutative association scheme such that $|S| > 3$. Then (X, S) contains exactly two nonlinear irreducible characters if and only if one of the following holds:*

(i) $|\mathrm{O}^{\vartheta}(S)| = 3$ *and S is the wreath product of association schemes $(X, S)_{x\mathrm{O}^{\vartheta}(S)}$ and $(X/\!\!/\mathrm{O}^{\vartheta}(S), S/\!\!/\mathrm{O}^{\vartheta}(S))$ for $x \in X$,*

(ii) $|\mathrm{O}^{\vartheta}(S)| = 2$ *and there exists a strongly normal closed subset H of S such that $|H| = 4$, $|H : \mathrm{O}^{\vartheta}(S)| = 2$ and $\varphi(\sigma_s) = 0$ for every nonlinear irreducible character φ of S and every $s \in S \setminus H$.*

Proof. Let χ and ψ be two nonlinear irreducible characters of S. Put $T = \mathrm{O}^{\vartheta}(S)$. First we assume that $\chi_T \neq \psi_T$. It follows that $\mathrm{Irr}(T) = \{1_T, \chi_T, \psi_T\}$. From [4, Theorem 3.4], $\mathrm{Irr}(S/\!\!/T)$ acts on $\mathrm{Irr}(S)$ by

$$\varphi\zeta(\sigma_s) = \varphi(\sigma_s)\zeta(\sigma_{s^T}),$$

for every $\varphi \in \mathrm{Irr}(S)$ and $\zeta \in \mathrm{Irr}(S/\!\!/T)$. Since $\chi_T \neq \psi_T$, it follows that the orbits of χ and ψ have length 1. So for every $\zeta \in \mathrm{Irr}(S/\!\!/T)$ we have $\chi\zeta = \chi$ and $\psi\zeta = \psi$. Since for every $s \in S \setminus T$, there exists $\zeta \in \mathrm{Irr}(S/\!\!/T)$ such that $\zeta(\sigma_{s^T}) \neq 1$, from equalities $\chi(\sigma_s)\zeta(\sigma_{s^T}) = \chi(\sigma_s)$ and $\psi(\sigma_s)\zeta(\sigma_{s^T}) = \psi(\sigma_s)$, we conclude that $\chi(\sigma_s) = \psi(\sigma_s) = 0$,

for every $s \in S \setminus T$. Now we show that for every $s, t \in S \setminus T$, $s^T \neq t^T$ and hence $|S/\!\!/T| = |S| - |T| + 1$. Suppose that $s^T = t^T$. Then for every $\varphi \in \mathrm{Irr}(S/\!\!/T)$, we have

$$\varphi(\sigma_s)/n_s = \varphi(\sigma_{s^T}) = \varphi(\sigma_{t^T}) = \varphi(\sigma_t)/n_t.$$

Since $\chi(\sigma_s) = \psi(\sigma_t) = 0$, it follows that for every $\varphi \in \mathrm{Irr}(S)$, $\varphi(\sigma_s)/n_s = \varphi(\sigma_t)/n_t$. This is a contradiction since the character table of $\mathbb{C}S$ is a nonsingular matrix. So $|S/\!\!/T| = |S| - |T| + 1$. This implies that $|S| = |S/\!\!/T| + 2 = |S/\!\!/T| + |T| - 1$ and Lemma 2.5 shows that S is the wreath product of association schemes $(X, S)_{xT}$ and $(X/\!\!/T, S/\!\!/T)$.

Now we can assume that $\chi_T = \psi_T$. We show that $|T| = 2$. Suppose that $s, t \in T \setminus \{1\}$. Since for every $\varphi \in \mathrm{Irr}(S/\!\!/T)$, $\varphi(\sigma_s)/n_s = \varphi(\sigma_t)/n_t = 1$ and $\chi(\sigma_s) = \psi(\sigma_t)$, it follows that $s = t$. Hence $T = \{1, s\}$, for some $s \in T$. Then since $|S| = |S/\!\!/T| + 2$, it follows that there exist exactly two relations $g, h \in S \setminus T$ such that $g^T = h^T$. We show that $H = \{1, s, g, h\}$ is a strongly normal closed subset of S such that $|H : T| = 2$. To do this, first note that g and h are symmetric relations. Moreover, since $g^T = h^T$, we have $\sigma_g \sigma_h = a\sigma_s$, for some positive integer a. If for some relation f we have $f \in sg$, then since $\lambda_{sgf} n_f = \lambda_{fgs} n_s$, we conclude that $s \in fg$ and hence $f^T = g^T$. Our assumption on g and h shows that $f \in \{g, h\}$ and then $sg \subseteq \{g, h\}$. Similarly, $sh \subseteq \{g, h\}$. This implies that $H = \{1, s, g, h\}$ is a closed subset of S. Since $H/\!\!/T$ is a subgroup of the abelian group $S/\!\!/T$, it follows that H is also strongly normal closed subset of S. Clearly if H is thin, then $|H : T| = 2$. Otherwise, since χ_H and ψ_H are two nonlinear irreducible characters of H, we conclude that $|H : T| = 2$. Finally, we show that $\chi(\sigma_s) = \psi(\sigma_s) = 0$, for every $s \in S \setminus H$. To do so, we first note that $\mathrm{Irr}(S/\!\!/H)$ acts on $\mathrm{Irr}(S)$ by

$$\varphi\zeta(\sigma_s) = \varphi(\sigma_s)\zeta(\sigma_{s^H}),$$

for every $\varphi \in \mathrm{Irr}(S)$ and $\zeta \in \mathrm{Irr}(S/\!\!/H)$ (see [4, Theorem 3.4]). Since $\chi_H \neq \psi_H$ it follows that the orbits of χ and ψ have length 1. Then for every $\zeta \in \mathrm{Irr}(S/\!\!/H)$ we get $\chi\zeta = \chi$ and $\psi\zeta = \psi$. Since for every $s \in S \setminus H$, there exists $\zeta \in \mathrm{Irr}(S/\!\!/H)$ such that $\zeta(\sigma_{s^H}) \neq 1$, from equalities $\chi(\sigma_s)\zeta(\sigma_{s^H}) = \chi(\sigma_s)$ and $\psi(\sigma_s)\zeta(\sigma_{s^H}) = \psi(\sigma_s)$, we conclude that $\chi(\sigma_s) = \psi(\sigma_s) = 0$, for every $s \in S \setminus H$.

Conversely, if (X, S) satisfying condition (i), then it follows from Theorem 2.7 that (X, S) contains exactly two nonlinear irreducible characters. So suppose (X, S) satisfies (ii). Since $|H| = 4$ and $|H : \mathrm{O}_\vartheta(S)| = 2$, we have exactly two irreducible characters $\lambda, \mu \in \mathrm{Irr}(H) - \mathrm{Irr}(H/\!\!/\mathrm{O}_\vartheta(S))$. Then for every nonlinear irreducible character $\varphi \in \mathrm{Irr}(S)$, $\varphi_H \in \{\lambda, \mu\}$. Moreover, if $\varphi, \phi \in \mathrm{Irr}(S)$ are two nonlinear characters such that $\varphi_H = \phi_H$, then $\varphi = \phi$; indeed, $\varphi(\sigma_s) = \phi(\sigma_s) = 0$, for every $s \in S \setminus H$. This implies that (X, S) exactly contains two nonlinear irreducible characters λ^S and μ^S. The proof is now complete.

■

From the above theorem we can obtain the following characterization of finite groups with two nonlinear irreducible characters.

Corollary 3.7. *A finite group G with at least three conjugacy classes has exactly two nonlinear irreducible characters if and only if one of the following holds:*

(i) *G' contains three conjugacy classes and for every $g \in G \setminus G'$, coset gG' is the conjugacy class of G containing g,*

(ii) *G' contains two conjugacy classes and there exists a normal subgroup H of G such that $G' \subseteq H$, H contains four conjugacy classes, $|H : G'| = 2$ and $\varphi(g) = 0$ for every nonlinear irreducible character φ and $g \in G \setminus H$.*

Proof. Let (G, S) be the group association scheme of G. Suppose $\{C_0 = \{1\}, \ldots, C_d\}$ is the conjugacy classes of G. For every conjugacy class C_i, assume $K_i = \sum_{g \in C_i} g$. First assume that G has two nonlinear irreducible characters. Then (G, S) also has two nonlinear irreducible characters and it follows from Theorem 3.6 that one of the following holds:

(1) $|\mathrm{O}^\vartheta(S)| = 3$ and S is the wreath product of association schemes $(G, S)_{g\mathrm{O}^\vartheta(S)}$ and $(G/\!\!/\mathrm{O}^\vartheta(S), S/\!\!/\mathrm{O}^\vartheta(S))$ for $g \in G$,

(2) $|\mathrm{O}^\vartheta(S)| = 2$ and there exists a strongly normal closed subset H of S such that $|H| = 4$, $|H : \mathrm{O}^\vartheta(S)| = 2$ and $\varphi(\sigma_s) = 0$ for every nonlinear irreducible character φ of S and every $s \in S \setminus H$.

It follows from (1) that G' contains three conjugacy classes C_0, C_1 and C_2 and for every conjugacy class C_i, Lemma 2.5 shows that $K_iK_1 = K_1K_i = |C_1|K_i$ and $K_iK_2 = K_2K_i = |C_2|K_i$. This yields $C_1C_i = C_i$ and $C_2C_i = C_i$. Then for every $x \in C_i$ we have $xG' \subseteq C_i$. Since from Lemma 2.6 we have $|xG'| = |G'| = |C_i|$ we obtain $xG' = C_i$ and statement (i) holds. Statement (ii) is immediate from (2).

Conversely, from statement (i) we have $|\mathrm{O}^\vartheta(S)| = 3$ and for every conjugacy class C_i of G and $x \in C_i$, $xC_1 = C_i$ and $xC_2 = C_i$. This implies that $C_1C_i = C_i$ and $C_2C_i = C_i$. So $K_iK_1 = K_1K_i = |C_1|K_i$ and $K_iK_2 = K_2K_i = |C_2|K_i$. Then Lemma 2.5 shows that S is the wreath product of association schemes $(G, S)_{g\mathrm{O}^\vartheta(S)}$ and $(G/\!\!/\mathrm{O}^\vartheta(S), S/\!\!/\mathrm{O}^\vartheta(S))$ for $g \in G$. Hence Theorem 3.6 yields (G, S) contains two nonlinear irreducible characters and so G also contains two nonlinear irreducible characters. If statement (ii) holds, then Theorem 3.6 shows that (G, S) contains two nonlinear irreducible characters and so G also contains two nonlinear irreducible characters.

■

The following theorem gives a class of association schemes with exactly two nonlinear irreducible characters.

Theorem 3.8. *Let (X, S) be an association scheme such that $|\mathrm{Irr}(S)| > 3$ and S' is symmetric. Then (X, S) contains exactly two nonlinear irreducible characters if and only if (X, S) is a group-like scheme and one of the following holds:*

(i) *$|\widetilde{S}'| = 3$ and $\widetilde{S}$ is the wreath product of association schemes $(X, S)_{x\widetilde{S}'}$ and $(X/\!\!/\widetilde{S}', \widetilde{S}/\!\!/\widetilde{S}')$ for $x \in X$,*

(ii) $|\widetilde{S'}| = 2$ *and there exists a strongly normal closed subset* H *of* S *such that* $|\widetilde{H}| = 4$, $|H : S'| = 2$ *and for every* $s \in S \setminus H$, $\chi(\sigma_s) = \psi(\sigma_s) = 0$.

Proof. First suppose (X, S) contains exactly two nonlinear irreducible characters. and let χ and ψ be two nonlinear irreducible characters of S. Since $S/\!\!/S'$ is an abelian group, $\mathrm{Irr}(S/\!\!/S')$ acts on $\mathrm{Irr}(S)$ by

$$\psi\zeta(\sigma_s) = \psi(\sigma_s)\zeta(\sigma_{s^{S'}})$$

for every $\psi \in \mathrm{Irr}(S)$ and $\zeta \in \mathrm{Irr}(S/\!\!/S')$; (see [4, Theorem 3.4]). Since χ and ψ are only nonlinear irreducible characters of S, it follows that the orbits of χ and ψ have length 1 or χ and ψ lie in the same orbit.

First we assume that the orbits of χ and ψ have length 1. This implies that $\chi_{S'} \neq \psi_{S'}$ and for every $\zeta \in \mathrm{Irr}(S/\!\!/S')$, $\chi\zeta = \chi$ and $\psi\zeta = \psi$. Since for every $s \in S \setminus S'$, there exists $\zeta \in \mathrm{Irr}(S/\!\!/S')$ such that $\zeta(\sigma_{s^{S'}}) \neq 1$, from equalities $\chi(\sigma_s) = \chi(\sigma_s)\zeta(\sigma_{s^{S'}})$ and $\psi(\sigma_s) = \psi(\sigma_s)\zeta(\sigma_{s^{S'}})$ we have $\chi(\sigma_s) = \psi(\sigma_s) = 0$. Now we prove that (X, S) is a group-like scheme. To do so, we show that for every $\lambda, \mu \in \mathrm{Irr}(S)$, $\lambda\mu(\sigma_s\sigma_h) = \lambda\mu(\sigma_h\sigma_s)$ for every $s, h \in S$. Then from Lemma 2.9 (X, S) is a group-like scheme. Let $\lambda, \mu \in \mathrm{Irr}(S)$. Clearly, if λ or μ belongs to $\mathrm{Irr}(S/\!\!/S')$, then $\lambda\mu \in \mathrm{Irr}(S)$ and the result follows. So we can assume that $\lambda, \mu \in \mathrm{Irr}(S) - \mathrm{Irr}(S/\!\!/S')$. Then $\lambda, \mu \in \{\chi, \psi\}$. We show that $\lambda\mu(\sigma_s\sigma_h) = \lambda\mu(\sigma_h\sigma_s)$ for every $s, h \in S$. Clearly, if $s, h \in S'$, then $\sigma_s\sigma_h = \sigma_h\sigma_s$ and so $\lambda\mu(\sigma_s\sigma_h) = \lambda\mu(\sigma_h\sigma_s)$. Moreover, if $s \in S'$ and $h \in S \setminus S'$ or $s, h \in S \setminus S'$ and $h \neq s^*$, then

$$\sigma_s\sigma_h \sum_{k \in S \setminus S'} a_k\sigma_k,$$

and so

$$\lambda\mu(\sigma_s\sigma_h) = \sum_{k \in S \setminus S'} a_k\lambda\mu(\sigma_k) = \sum_{k \in S \setminus S'} a_k \frac{\lambda(\sigma_k)\mu(\sigma_h)}{n_k} = 0$$

indeed, $\lambda(\sigma_s) = \mu(\sigma_s) = 0$, for every $s \in S \setminus S'$. Similarly, $\lambda\mu(\sigma_h\sigma_s) = 0$. Finally, we can assume that $s, h \in S \setminus S'$ and $h = s^*$. Then since S' is strongly normal and symmetric we have

$$\sigma_s\sigma_{s^*} = \sum_{k \in S'} a_k\sigma_k = \sum_{k \in S'} a_k\sigma_{k^*} = \sigma_{s^*}\sigma_s$$

and so $\lambda\mu(\sigma_s\sigma_{s^*}) = \lambda\mu(\sigma_{s^*}\sigma_s)$. Hence (X, S) is a group-like scheme.

Now consider association scheme $(X, \widetilde{S})$ and put $\widetilde{\chi} = \frac{\chi}{\chi(1)}$ and $\widetilde{\psi} = \frac{\psi}{\psi(1)}$. Then $\widetilde{\chi}$ and $\widetilde{\psi}$ are nonlinear irreducible characters of $\widetilde{S}$; see [2]. Since $\widetilde{\chi}(\sigma_{\widetilde{s}}) \neq \widetilde{\psi}(\sigma_{\widetilde{s}})$, for every $\widetilde{s} \in \widetilde{S'}$, statement (i) follows from part (i) of Theorem 3.6.

Now we assume that χ and ψ lie in the same orbit. Then $\chi_{S'} = \psi_{S'}$ and $\{\chi, \psi\}$ forms an orbit of length 2. So $S/\!\!/S'$ has a subgroup of order 2. Then there exists a strongly normal closed subset H of S such that $S' \subseteq H$ and $|H : S'| = 2$.

Clearly, $\chi_H \neq \psi_H$. We show that H is commutative. To do so, we prove that every irreducible character of H has degree 1. Suppose on contrary; that there exists $\varphi \in \mathrm{Irr}(H)$ such that $\varphi(1) > 1$. Since S' is commutative, it follows from Theorem 2.4 that $\varphi_{S'}$ is a sum of at most two distinct irreducible characters of S' and $\varphi = \lambda^H$ for some irreducible constituent λ of $\varphi_{S'}$. On the other hand, since $\chi_{S'} = \psi_{S'}$, we conclude that $\lambda^S = a\chi + b\psi$ for some integers a and b. Then

$$a\chi + b\psi = \lambda^S = (\lambda^H)^S = \varphi^S.$$

This implies that $(\chi_H, \varphi) > 0$ and $(\psi_H, \varphi) > 0$ and then it follows from Theorem 2.3 that

$$\chi_H = \psi_H = e\sum_{i=1}^{n} \varphi_i,$$

where $\varphi_i, 1 \leq i \leq n$, are $S/\!\!/H$-conjugate irreducible characters of φ. This is a contradiction, indeed $\chi_H \neq \psi_H$. Hence H is commutative.

Now by a similar way as above, one can see that for every $\lambda, \mu \in \mathrm{Irr}(S)$, $\lambda\mu(\sigma_s\sigma_h) = \lambda\mu(\sigma_h\sigma_s)$ for every $s, h \in S$. Note that S' is symmetric, H is commutative and $\chi(\sigma_s) = \psi(\sigma_s) = 0$ for every $s \in S \setminus H$. So it follows from Lemma 2.9 that (X, S) is a group-like scheme. Moreover, since association scheme $(X, \widetilde{S})$ has exactly two nonlinear irreducible character, it follows from statement (ii) of Theorem 3.6 that $|\mathrm{O}^{\vartheta}(\widetilde{S})| = |\widetilde{S'}| = 2$ and $|\widetilde{H}| = 4$.

Conversely, it follows from Theorem 3.6 that $(X, \widetilde{S})$ has two nonlinear irreducible character. Since $m_{\widetilde{\chi}} = \chi(1)m_\chi$, for every $\chi \in \mathrm{Irr}(S)$ we conclude that (X, S) has two nonlinear irreducible character. The proof is now complete.

■

Remark 3.9. *The conditions* $|\mathrm{Irr}(S)| > 3$ *and symmetry of* S' *in Theorem 3.8 are necessary conditions; see example below.*

Example 3.10. *Let* (X, S) *be the association scheme of order* 21*, No.* 19 *in [7], where* $S = \{s_0, \cdots, s_6\}$*. Then from [7] the character table of the adjacency algebra of* S *is as follows.*

	σ_{s_0}	σ_{s_1}	σ_{s_2}	σ_{s_3}	σ_{s_4}	σ_{s_5}	m_χ
χ_1	1	2	2	4	4	8	1
χ_2	1	−1	−1	1	1	−1	8
χ_3	2	1	1	−2	−2	0	6

One can see that (X, S) *has two nonlinear irreducible characters, but it is not a group-like scheme and the assertion of Theorem 3.8 does not hold for* (X, S)*. Moreover, if we consider the association scheme* $H = S\wr K$*, where* K *is the trivial scheme of order* 2*, then* H *has order* 42 *and from Theorem 2.7, the character table of* H *is*

as follows.

	σ_{s_0}	σ_{s_1}	σ_{s_2}	σ_{s_3}	σ_{s_4}	σ_{s_5}	σ_{s_6}	m_χ
χ_1	1	2	2	4	4	8	21	1
χ_2	1	2	2	4	4	8	-21	1
χ_2	1	-1	-1	1	1	-1	0	16
χ_3	2	1	1	-2	-2	0	0	12

It is easy to see that H is not a group-like scheme. Although, H contains exactly two nonlinear irreducible characters, but H' is not symmetric and so the conclusion of Theorem 3.8 does not hold for H.

References

[1] A. Hanaki, Clifford theory for association schemes, J. Algebra, 321 (2009) 1686-1695.

[2] A. Hanaki, Nilpotent schemes and group-like schemes, J. Combin. Theory Ser. A, 115 (2008) 226-236.

[3] A. Hanaki, Characters of association schemes containing a strongly normal closed subset of prime index, Proc. Amer. Math. Soc., 135 (2007) 2683-2687.

[4] A. Hanaki, Character products of association schemes, J. Algebra 283 (2005) 596-603.

[5] A. Hanaki, Representations of association schemes and their factor schemes, Graphs Combin., 19 (2003) 195-201.

[6] A. Hanaki, K. Hirotsuka, Irreducible representations of wreath products of association schemes, J. Algebraic Combin., 18 (2003) 47-52.

[7] A. Hanaki, Data for association schemes. http://kissme.shinshu-u.ac.jp/as/data.

[8] M. Hirasaka, Kyoung-tark Kim, On p-covalenced association schemes, J. Combin. Theory Ser. A, 118 (2011) 1-8.

[9] A.S. Muktibodh, Con-Cos groups, Review Bulletin of Calcutta Mathematical Society, 11 (2003), 53-56.

[10] B. Xu, On wreath products of C-algebras, table algebras, and association schemes, J. Algebra, 344 (2011) 268-283.

[11] P.-H. Zieschang, An Algebraic Approach to Association Schemes, Lecture Notes in Math., vol. 1628, Springer, Berlin, 1996.

[12] P.-H. Zieschang, Theory of Association Schemes, Springer Monographs in Mathematics, Springer-Berlin, 2005.